

## EXERCISES 3.4

## Derivatives

In Exercises 1–12, find  $dy/dx$ .

1.  $y = -10x + 3 \cos x$
2.  $y = \frac{3}{x} + 5 \sin x$
3.  $y = \csc x - 4\sqrt{x} + 7$
4.  $y = x^2 \cot x - \frac{1}{x^2}$
5.  $y = (\sec x + \tan x)(\sec x - \tan x)$
6.  $y = (\sin x + \cos x) \sec x$
7.  $y = \frac{\cot x}{1 + \cot x}$
8.  $y = \frac{\cos x}{1 + \sin x}$
9.  $y = \frac{4}{\cos x} + \frac{1}{\tan x}$
10.  $y = \frac{\cos x}{x} + \frac{x}{\cos x}$
11.  $y = x^2 \sin x + 2x \cos x - 2 \sin x$
12.  $y = x^2 \cos x - 2x \sin x - 2 \cos x$

In Exercises 13–16, find  $ds/dt$ .

13.  $s = \tan t - t$
14.  $s = t^2 - \sec t + 1$
15.  $s = \frac{1 + \csc t}{1 - \csc t}$
16.  $s = \frac{\sin t}{1 - \cos t}$

In Exercises 17–20, find  $dr/d\theta$ .

17.  $r = 4 - \theta^2 \sin \theta$
18.  $r = \theta \sin \theta + \cos \theta$
19.  $r = \sec \theta \csc \theta$
20.  $r = (1 + \sec \theta) \sin \theta$

In Exercises 21–24, find  $dp/dq$ .

21.  $p = 5 + \frac{1}{\cot q}$
22.  $p = (1 + \csc q) \cos q$
23.  $p = \frac{\sin q + \cos q}{\cos q}$
24.  $p = \frac{\tan q}{1 + \tan q}$

25. Find  $y''$  if

- a.  $y = \csc x$ .
- b.  $y = \sec x$ .

26. Find  $y^{(4)} = d^4 y/dx^4$  if

- a.  $y = -2 \sin x$ .
- b.  $y = 9 \cos x$ .

## Tangent Lines



In Exercises 27–30, graph the curves over the given intervals, together with their tangents at the given values of  $x$ . Label each curve and tangent with its equation.

27.  $y = \sin x$ ,  $-3\pi/2 \leq x \leq 2\pi$   
 $x = -\pi, 0, 3\pi/2$
28.  $y = \tan x$ ,  $-\pi/2 < x < \pi/2$   
 $x = -\pi/3, 0, \pi/3$
29.  $y = \sec x$ ,  $-\pi/2 < x < \pi/2$   
 $x = -\pi/3, \pi/4$
30.  $y = 1 + \cos x$ ,  $-3\pi/2 \leq x \leq 2\pi$   
 $x = -\pi/3, 3\pi/2$

**T** Do the graphs of the functions in Exercises 31–34 have any horizontal tangents in the interval  $0 \leq x \leq 2\pi$ ? If so, where? If not, why not? Visualize your findings by graphing the functions with a grapher.

31.  $y = x + \sin x$
32.  $y = 2x + \sin x$
33.  $y = x - \cot x$
34.  $y = x + 2 \cos x$
35. Find all points on the curve  $y = \tan x$ ,  $-\pi/2 < x < \pi/2$ , where the tangent line is parallel to the line  $y = 2x$ . Sketch the curve and tangent(s) together, labeling each with its equation.
36. Find all points on the curve  $y = \cot x$ ,  $0 < x < \pi$ , where the tangent line is parallel to the line  $y = -x$ . Sketch the curve and tangent(s) together, labeling each with its equation.

In Exercises 37 and 38, find an equation for (a) the tangent to the curve at  $P$  and (b) the horizontal tangent to the curve at  $Q$ .



## Trigonometric Limits

Find the limits in Exercises 39–44.

39.  $\lim_{x \rightarrow 2} \sin\left(\frac{1}{x} - \frac{1}{2}\right)$

40.  $\lim_{x \rightarrow -\pi/6} \sqrt{1 + \cos(\pi \csc x)}$

41.  $\lim_{x \rightarrow 0} \sec\left[\cos x + \pi \tan\left(\frac{\pi}{4 \sec x}\right) - 1\right]$

42.  $\lim_{x \rightarrow 0} \sin\left(\frac{\pi + \tan x}{\tan x - 2 \sec x}\right)$

43.  $\lim_{t \rightarrow 0} \tan\left(1 - \frac{\sin t}{t}\right)$

44.  $\lim_{\theta \rightarrow 0} \cos\left(\frac{\pi \theta}{\sin \theta}\right)$

## Simple Harmonic Motion

The equations in Exercises 45 and 46 give the position  $s = f(t)$  of a body moving on a coordinate line ( $s$  in meters,  $t$  in seconds). Find the body's velocity, speed, acceleration, and jerk at time  $t = \pi/4$  sec.

45.  $s = 2 - 2 \sin t$

46.  $s = \sin t + \cos t$

## Theory and Examples

47. Is there a value of  $c$  that will make

$$f(x) = \begin{cases} \frac{\sin^2 3x}{x^2}, & x \neq 0 \\ c, & x = 0 \end{cases}$$

continuous at  $x = 0$ ? Give reasons for your answer.

48. Is there a value of  $b$  that will make

$$g(x) = \begin{cases} x + b, & x < 0 \\ \cos x, & x \geq 0 \end{cases}$$

continuous at  $x = 0$ ? Differentiable at  $x = 0$ ? Give reasons for your answers.

49. Find  $d^{999}/dx^{999}(\cos x)$ .

50. Derive the formula for the derivative with respect to  $x$  of

a.  $\sec x$ .    b.  $\csc x$ .    c.  $\cot x$ .

**T** 51. Graph  $y = \cos x$  for  $-\pi \leq x \leq 2\pi$ . On the same screen, graph

$$y = \frac{\sin(x+h) - \sin x}{h}$$

for  $h = 1, 0.5, 0.3$ , and  $0.1$ . Then, in a new window, try  $h = -1, -0.5$ , and  $-0.3$ . What happens as  $h \rightarrow 0^+$ ? As  $h \rightarrow 0^-$ ? What phenomenon is being illustrated here?

**T** 52. Graph  $y = -\sin x$  for  $-\pi \leq x \leq 2\pi$ . On the same screen, graph

$$y = \frac{\cos(x+h) - \cos x}{h}$$

for  $h = 1, 0.5, 0.3$ , and  $0.1$ . Then, in a new window, try  $h = -1, -0.5$ , and  $-0.3$ . What happens as  $h \rightarrow 0^+$ ? As  $h \rightarrow 0^-$ ? What phenomenon is being illustrated here?


**T** 53. **Centered difference quotients** The *centered difference quotient*

$$\frac{f(x+h) - f(x-h)}{2h}$$

is used to approximate  $f'(x)$  in numerical work because (1) its limit as  $h \rightarrow 0$  equals  $f'(x)$  when  $f'(x)$  exists, and (2) it usually gives a better approximation of  $f'(x)$  for a given value of  $h$  than Fermat's difference quotient

$$\frac{f(x+h) - f(x)}{h}.$$

See the accompanying figure.



- a. To see how rapidly the centered difference quotient for  $f(x) = \sin x$  converges to  $f'(x) = \cos x$ , graph  $y = \cos x$  together with

$$y = \frac{\sin(x+h) - \sin(x-h)}{2h}$$

over the interval  $[-\pi, 2\pi]$  for  $h = 1, 0.5$ , and  $0.3$ . Compare the results with those obtained in Exercise 51 for the same values of  $h$ .

- b. To see how rapidly the centered difference quotient for  $f(x) = \cos x$  converges to  $f'(x) = -\sin x$ , graph  $y = -\sin x$  together with

$$y = \frac{\cos(x+h) - \cos(x-h)}{2h}$$

over the interval  $[-\pi, 2\pi]$  for  $h = 1, 0.5$ , and  $0.3$ . Compare the results with those obtained in Exercise 52 for the same values of  $h$ .

54. **A caution about centered difference quotients** (Continuation of Exercise 53.) The quotient

$$\frac{f(x+h) - f(x-h)}{2h}$$

may have a limit as  $h \rightarrow 0$  when  $f$  has no derivative at  $x$ . As a case in point, take  $f(x) = |x|$  and calculate

$$\lim_{h \rightarrow 0} \frac{|0+h| - |0-h|}{2h}.$$

As you will see, the limit exists even though  $f(x) = |x|$  has no derivative at  $x = 0$ . *Moral:* Before using a centered difference quotient, be sure the derivative exists.

- T 55. Slopes on the graph of the tangent function** Graph  $y = \tan x$  and its derivative together on  $(-\pi/2, \pi/2)$ . Does the graph of the tangent function appear to have a smallest slope? a largest slope? Is the slope ever negative? Give reasons for your answers.

- T 56. Slopes on the graph of the cotangent function** Graph  $y = \cot x$  and its derivative together for  $0 < x < \pi$ . Does the graph of the cotangent function appear to have a smallest slope? A largest slope? Is the slope ever positive? Give reasons for your answers.

- T 57. Exploring  $(\sin kx)/x$**  Graph  $y = (\sin x)/x$ ,  $y = (\sin 2x)/x$ , and  $y = (\sin 4x)/x$  together over the interval  $-2 \leq x \leq 2$ . Where does each graph appear to cross the  $y$ -axis? Do the graphs really intersect the axis? What would you expect the graphs of  $y = (\sin 5x)/x$  and  $y = (\sin(-3x))/x$  to do as  $x \rightarrow 0$ ? Why? What about the graph of  $y = (\sin kx)/x$  for other values of  $k$ ? Give reasons for your answers.

- T 58. Radians versus degrees: degree mode derivatives** What happens to the derivatives of  $\sin x$  and  $\cos x$  if  $x$  is measured in degrees instead of radians? To find out, take the following steps.

- a. With your graphing calculator or computer grapher in *degree mode*, graph

$$f(h) = \frac{\sin h}{h}$$

and estimate  $\lim_{h \rightarrow 0} f(h)$ . Compare your estimate with  $\pi/180$ . Is there any reason to believe the limit *should* be  $\pi/180$ ?

- b. With your grapher still in degree mode, estimate

$$\lim_{h \rightarrow 0} \frac{\cos h - 1}{h}.$$

- c. Now go back to the derivation of the formula for the derivative of  $\sin x$  in the text and carry out the steps of the derivation using degree-mode limits. What formula do you obtain for the derivative?
- d. Work through the derivation of the formula for the derivative of  $\cos x$  using degree-mode limits. What formula do you obtain for the derivative?
- e. The disadvantages of the degree-mode formulas become apparent as you start taking derivatives of higher order. Try it. What are the second and third degree-mode derivatives of  $\sin x$  and  $\cos x$ ?