



Chapter 5 Practice Exercises

Finite Sums and Estimates

1. The accompanying figure shows the graph of the velocity (ft/sec) of a model rocket for the first 8 sec after launch. The rocket accelerated straight up for the first 2 sec and then coasted to reach its maximum height at $t = 8$ sec.



- a. Assuming that the rocket was launched from ground level, about how high did it go? (This is the rocket in Section 3.3, Exercise 17, but you do not need to do Exercise 17 to do the exercise here.)
- b. Sketch a graph of the rocket's height aboveground as a function of time for $0 \leq t \leq 8$.
2. a. The accompanying figure shows the velocity (m/sec) of a body moving along the s -axis during the time interval from $t = 0$ to $t = 10$ sec. About how far did the body travel during those 10 sec?
- b. Sketch a graph of s as a function of t for $0 \leq t \leq 10$ assuming $s(0) = 0$.



3. Suppose that $\sum_{k=1}^{10} a_k = -2$ and $\sum_{k=1}^{10} b_k = 25$. Find the value of

a. $\sum_{k=1}^{10} \frac{a_k}{4}$ b. $\sum_{k=1}^{10} (b_k - 3a_k)$

c. $\sum_{k=1}^{10} (a_k + b_k - 1)$ d. $\sum_{k=1}^{10} \left(\frac{5}{2} - b_k \right)$

4. Suppose that $\sum_{k=1}^{20} a_k = 0$ and $\sum_{k=1}^{20} b_k = 7$. Find the values of

a. $\sum_{k=1}^{20} 3a_k$ b. $\sum_{k=1}^{20} (a_k + b_k)$

c. $\sum_{k=1}^{20} \left(\frac{1}{2} - \frac{2b_k}{7} \right)$ d. $\sum_{k=1}^{20} (a_k - 2)$

Definite Integrals

In Exercises 5–8, express each limit as a definite integral. Then evaluate the integral to find the value of the limit. In each case, P is a partition of the given interval and the numbers c_k are chosen from the subintervals of P .

5. $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (2c_k - 1)^{-1/2} \Delta x_k$, where P is a partition of $[1, 5]$
6. $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n c_k (c_k^2 - 1)^{1/3} \Delta x_k$, where P is a partition of $[1, 3]$
7. $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n \left(\cos \left(\frac{c_k}{2} \right) \right) \Delta x_k$, where P is a partition of $[-\pi, 0]$
8. $\lim_{\|P\| \rightarrow 0} \sum_{k=1}^n (\sin c_k)(\cos c_k) \Delta x_k$, where P is a partition of $[0, \pi/2]$
9. If $\int_{-2}^2 3f(x) dx = 12$, $\int_{-2}^5 f(x) dx = 6$, and $\int_{-2}^5 g(x) dx = 2$, find the values of the following.
- a. $\int_{-2}^2 f(x) dx$ b. $\int_2^5 f(x) dx$
- c. $\int_5^{-2} g(x) dx$ d. $\int_{-2}^5 (-\pi g(x)) dx$
- e. $\int_{-2}^5 \left(\frac{f(x) + g(x)}{5} \right) dx$
10. If $\int_0^2 f(x) dx = \pi$, $\int_0^2 7g(x) dx = 7$, and $\int_0^1 g(x) dx = 2$, find the values of the following.

a. $\int_0^2 g(x) dx$ b. $\int_1^2 g(x) dx$

c. $\int_2^0 f(x) dx$ d. $\int_0^2 \sqrt{2} f(x) dx$

e. $\int_0^2 (g(x) - 3f(x)) dx$

Area

In Exercise 11–14, find the total area of the region between the graph of f and the x -axis.

11. $f(x) = x^2 - 4x + 3$, $0 \leq x \leq 3$
12. $f(x) = 1 - (x^2/4)$, $-2 \leq x \leq 3$

13. $f(x) = 5 - 5x^{2/3}, \quad -1 \leq x \leq 8$


14. $f(x) = 1 - \sqrt{x}, \quad 0 \leq x \leq 4$

Find the areas of the regions enclosed by the curves and lines in Exercises 15–26.


15. $y = x, \quad y = 1/x^2, \quad x = 2$

16. $y = x, \quad y = 1/\sqrt{x}, \quad x = 2$

17. $\sqrt{x} + \sqrt{y} = 1, \quad x = 0, \quad y = 0$



18. $x^3 + \sqrt{y} = 1, \quad x = 0, \quad y = 0, \quad \text{for } 0 \leq x \leq 1$



19. $x = 2y^2, \quad x = 0, \quad y = 3$

20. $x = 4 - y^2, \quad x = 0$

21. $y^2 = 4x, \quad y = 4x - 2$

22. $y^2 = 4x + 4, \quad y = 4x - 16$

23. $y = \sin x, \quad y = x, \quad 0 \leq x \leq \pi/4$

24. $y = |\sin x|, \quad y = 1, \quad -\pi/2 \leq x \leq \pi/2$

25. $y = 2 \sin x, \quad y = \sin 2x, \quad 0 \leq x \leq \pi$

26. $y = 8 \cos x, \quad y = \sec^2 x, \quad -\pi/3 \leq x \leq \pi/3$

27. Find the area of the “triangular” region bounded on the left by $x + y = 2$, on the right by $y = x^2$, and above by $y = 2$.

28. Find the area of the “triangular” region bounded on the left by $y = \sqrt{x}$, on the right by $y = 6 - x$, and below by $y = 1$.

29. Find the extreme values of $f(x) = x^3 - 3x^2$ and find the area of the region enclosed by the graph of f and the x -axis.

30. Find the area of the region cut from the first quadrant by the curve $x^{1/2} + y^{1/2} = a^{1/2}$.

31. Find the total area of the region enclosed by the curve $x = y^{2/3}$ and the lines $x = y$ and $y = -1$.

32. Find the total area of the region between the curves $y = \sin x$ and $y = \cos x$ for $0 \leq x \leq 3\pi/2$.

Initial Value Problems

33. Show that $y = x^2 + \int_1^x \frac{1}{t} dt$ solves the initial value problem

$$\frac{d^2 y}{dx^2} = 2 - \frac{1}{x^2}; \quad y'(1) = 3, \quad y(1) = 1.$$

34. Show that $y = \int_0^x (1 + 2\sqrt{\sec t}) dt$ solves the initial value problem

$$\frac{d^2 y}{dx^2} = \sqrt{\sec x} \tan x; \quad y'(0) = 3, \quad y(0) = 0.$$

Express the solutions of the initial value problems in Exercises 35 and 36 in terms of integrals.

35. $\frac{dy}{dx} = \frac{\sin x}{x}, \quad y(5) = -3$

36. $\frac{dy}{dx} = \sqrt{2 - \sin^2 x}, \quad y(-1) = 2$

Evaluating Indefinite Integrals

Evaluate the integrals in Exercises 37–44.

37. $\int 2(\cos x)^{-1/2} \sin x dx$

38. $\int (\tan x)^{-3/2} \sec^2 x dx$

39. $\int (2\theta + 1 + 2 \cos(2\theta + 1)) d\theta$

40. $\int \left(\frac{1}{\sqrt{2\theta - \pi}} + 2 \sec^2(2\theta - \pi) \right) d\theta$

41. $\int \left(t - \frac{2}{t} \right) \left(t + \frac{2}{t} \right) dt$

42. $\int \frac{(t+1)^2 - 1}{t^4} dt$

43. $\int \sqrt{t} \sin(2t^{3/2}) dt$

44. $\int \sec \theta \tan \theta \sqrt{1 + \sec \theta} d\theta$

Evaluating Definite Integrals

Evaluate the integrals in Exercises 45–70.

45. $\int_{-1}^1 (3x^2 - 4x + 7) dx$

46. $\int_0^1 (8s^3 - 12s^2 + 5) ds$

47. $\int_1^2 \frac{4}{v^2} dv$

48. $\int_1^{27} x^{-4/3} dx$

49. $\int_1^4 \frac{dt}{t\sqrt{t}}$

50. $\int_1^4 \frac{(1 + \sqrt{u})^{1/2}}{\sqrt{u}} du$

51. $\int_0^1 \frac{36 dx}{(2x + 1)^3}$

52. $\int_0^1 \frac{dr}{\sqrt[3]{(7 - 5r)^2}}$

53. $\int_{1/8}^1 x^{-1/3} (1 - x^{2/3})^{3/2} dx$

54. $\int_0^{1/2} x^3 (1 + 9x^4)^{-3/2} dx$

55. $\int_0^{\pi} \sin^2 5r dr$

56. $\int_0^{\pi/4} \cos^2 \left(4t - \frac{\pi}{4} \right) dt$

57. $\int_0^{\pi/3} \sec^2 \theta \, d\theta$ 58. $\int_{\pi/4}^{3\pi/4} \csc^2 x \, dx$
59. $\int_{\pi}^{3\pi} \cot^2 \frac{x}{6} \, dx$ 60. $\int_0^{\pi} \tan^2 \frac{\theta}{3} \, d\theta$
61. $\int_{-\pi/3}^0 \sec x \tan x \, dx$ 62. $\int_{\pi/4}^{3\pi/4} \csc z \cot z \, dz$
63. $\int_0^{\pi/2} 5(\sin x)^{3/2} \cos x \, dx$ 64. $\int_{-1}^1 2x \sin(1 - x^2) \, dx$
65. $\int_{-\pi/2}^{\pi/2} 15 \sin^4 3x \cos 3x \, dx$ 66. $\int_0^{2\pi/3} \cos^{-4} \left(\frac{x}{2} \right) \sin \left(\frac{x}{2} \right) \, dx$
67. $\int_0^{\pi/2} \frac{3 \sin x \cos x}{\sqrt{1 + 3 \sin^2 x}} \, dx$ 68. $\int_0^{\pi/4} \frac{\sec^2 x}{(1 + 7 \tan x)^{2/3}} \, dx$
69. $\int_0^{\pi/3} \frac{\tan \theta}{\sqrt{2 \sec \theta}} \, d\theta$ 70. $\int_{\pi^2/36}^{\pi^2/4} \frac{\cos \sqrt{t}}{\sqrt{t \sin \sqrt{t}}} \, dt$

Average Values

71. Find the average value of $f(x) = mx + b$
- over $[-1, 1]$
 - over $[-k, k]$
72. Find the average value of
- $y = \sqrt{3x}$ over $[0, 3]$
 - $y = \sqrt{ax}$ over $[0, a]$
73. Let f be a function that is differentiable on $[a, b]$. In Chapter 2 we defined the average rate of change of f over $[a, b]$ to be

$$\frac{f(b) - f(a)}{b - a}$$

and the instantaneous rate of change of f at x to be $f'(x)$. In this chapter we defined the average value of a function. For the new definition of average to be consistent with the old one, we should have

$$\frac{f(b) - f(a)}{b - a} = \text{average value of } f' \text{ on } [a, b].$$

Is this the case? Give reasons for your answer.

74. Is it true that the average value of an integrable function over an interval of length 2 is half the function's integral over the interval? Give reasons for your answer.

- T** 75. Compute the average value of the temperature function

$$f(x) = 37 \sin \left(\frac{2\pi}{365} (x - 101) \right) + 25$$

for a 365-day year. This is one way to estimate the annual mean air temperature in Fairbanks, Alaska. The National Weather Service's official figure, a numerical average of the daily normal mean air temperatures for the year, is 25.7°F, which is slightly higher than the average value of $f(x)$. Figure 3.33 shows why.

- T** 76. **Specific heat of a gas** Specific heat C_v is the amount of heat required to raise the temperature of a given mass of gas with con-

stant volume by 1°C, measured in units of cal/deg-mole (calories per degree gram molecule). The specific heat of oxygen depends on its temperature T and satisfies the formula

$$C_v = 8.27 + 10^{-5} (26T - 1.87T^2).$$

Find the average value of C_v for $20^\circ \leq T \leq 675^\circ\text{C}$ and the temperature at which it is attained.


Differentiating Integrals

In Exercises 77–80, find dy/dx .

77. $y = \int_2^x \sqrt{2 + \cos^3 t} \, dt$ 78. $y = \int_2^{7x^2} \sqrt{2 + \cos^3 t} \, dt$
79. $y = \int_x^1 \frac{6}{3 + t^4} \, dt$ 80. $y = \int_{\sec x}^2 \frac{1}{t^2 + 1} \, dt$

Theory and Examples

81. Is it true that every function $y = f(x)$ that is differentiable on $[a, b]$ is itself the derivative of some function on $[a, b]$? Give reasons for your answer.
82. Suppose that $F(x)$ is an antiderivative of $f(x) = \sqrt{1 + x^4}$. Express $\int_0^1 \sqrt{1 + x^4} \, dx$ in terms of F and give a reason for your answer.
83. Find dy/dx if $y = \int_x^1 \sqrt{1 + t^2} \, dt$. Explain the main steps in your calculation.
84. Find dy/dx if $y = \int_{\cos x}^0 (1/(1 - t^2)) \, dt$. Explain the main steps in your calculation.
85. **A new parking lot** To meet the demand for parking, your town has allocated the area shown here. As the town engineer, you have been asked by the town council to find out if the lot can be built for \$10,000. The cost to clear the land will be \$0.10 a square foot, and the lot will cost \$2.00 a square foot to pave. Can the job be done for \$10,000? Use a lower sum estimate to see. (Answers may vary slightly, depending on the estimate used.)



86. Skydivers A and B are in a helicopter hovering at 6400 ft. Skydiver A jumps and descends for 4 sec before opening her parachute. The helicopter then climbs to 7000 ft and hovers there. Forty-five seconds after A leaves the aircraft, B jumps and descends for 13 sec before opening his parachute. Both skydivers descend at 16 ft/sec with parachutes open. Assume that the skydivers fall freely (no effective air resistance) before their parachutes open.
- At what altitude does A's parachute open?
 - At what altitude does B's parachute open?
 - Which skydiver lands first?

Average Daily Inventory

Average value is used in economics to study such things as average daily inventory. If $I(t)$ is the number of radios, tires, shoes, or whatever product a firm has on hand on day t (we call I an **inventory function**), the average value of I over a time period $[0, T]$ is called the firm's average daily inventory for the period.

$$\text{Average daily inventory} = \text{av}(I) = \frac{1}{T} \int_0^T I(t) dt.$$

If h is the dollar cost of holding one item per day, the product $\text{av}(I) \cdot h$ is the **average daily holding cost** for the period.

87. As a wholesaler, Tracey Burr Distributors receives a shipment of 1200 cases of chocolate bars every 30 days. TBD sells the chocolate to retailers at a steady rate, and t days after a shipment arrives, its inventory of cases on hand is $I(t) = 1200 - 40t$, $0 \leq t \leq 30$. What is TBD's average daily inventory for the 30-day period? What is its average daily holding cost if the cost of holding one case is 3¢ a day?
88. Rich Wholesale Foods, a manufacturer of cookies, stores its cases of cookies in an air-conditioned warehouse for shipment every 14 days. Rich tries to keep 600 cases on reserve to meet occasional peaks in demand, so a typical 14-day inventory function is $I(t) = 600 + 600t$, $0 \leq t \leq 14$. The daily holding cost for each case is 4¢ per day. Find Rich's average daily inventory and average daily holding cost.
89. Solon Container receives 450 drums of plastic pellets every 30 days. The inventory function (drums on hand as a function of days) is $I(t) = 450 - t^2/2$. Find the average daily inventory. If the holding cost for one drum is 2¢ per day, find the average daily holding cost.
90. Mitchell Mailorder receives a shipment of 600 cases of athletic socks every 60 days. The number of cases on hand t days after the shipment arrives is $I(t) = 600 - 20\sqrt{15t}$. Find the average daily inventory. If the holding cost for one case is 1/2¢ per day, find the average daily holding cost.