

EXERCISES 8.3

Expanding Quotients into Partial Fractions

Expand the quotients in Exercises 1–8 by partial fractions.

1.
$$\frac{5x - 13}{(x - 3)(x - 2)}$$

2.
$$\frac{5x - 7}{x^2 - 3x + 2}$$

3.
$$\frac{x + 4}{(x + 1)^2}$$

4.
$$\frac{2x + 2}{x^2 - 2x + 1}$$

5.
$$\frac{z + 1}{z^2(z - 1)}$$

6.
$$\frac{z}{z^3 - z^2 - 6z}$$

7.
$$\frac{t^2 + 8}{t^2 - 5t + 6}$$

8.
$$\frac{t^4 + 9}{t^4 + 9t^2}$$

Nonrepeated Linear Factors

In Exercises 9–16, express the integrands as a sum of partial fractions and evaluate the integrals.

9.
$$\int \frac{dx}{1 - x^2}$$

10.
$$\int \frac{dx}{x^2 + 2x}$$

11.
$$\int \frac{x + 4}{x^2 + 5x - 6} dx$$

12.
$$\int \frac{2x + 1}{x^2 - 7x + 12} dx$$

13.
$$\int_4^8 \frac{y dy}{y^2 - 2y - 3}$$

14.
$$\int_{1/2}^1 \frac{y + 4}{y^2 + y} dy$$

15.
$$\int \frac{dt}{t^3 + t^2 - 2t}$$

16.
$$\int \frac{x + 3}{2x^3 - 8x} dx$$

Repeated Linear Factors

In Exercises 17–20, express the integrands as a sum of partial fractions and evaluate the integrals.

17.
$$\int_0^1 \frac{x^3 dx}{x^2 + 2x + 1}$$

18.
$$\int_{-1}^0 \frac{x^3 dx}{x^2 - 2x + 1}$$

19.
$$\int \frac{dx}{(x^2 - 1)^2}$$

20.
$$\int \frac{x^2 dx}{(x - 1)(x^2 + 2x + 1)}$$

Irreducible Quadratic Factors

In Exercises 21–28, express the integrands as a sum of partial fractions and evaluate the integrals.

21.
$$\int_0^1 \frac{dx}{(x + 1)(x^2 + 1)}$$

22.
$$\int_1^{\sqrt{3}} \frac{3t^2 + t + 4}{t^3 + t} dt$$

23.
$$\int \frac{y^2 + 2y + 1}{(y^2 + 1)^2} dy$$

24.
$$\int \frac{8x^2 + 8x + 2}{(4x^2 + 1)^2} dx$$

25.
$$\int \frac{2s + 2}{(s^2 + 1)(s - 1)^3} ds$$

26.
$$\int \frac{s^4 + 81}{s(s^2 + 9)^2} ds$$

27.
$$\int \frac{2\theta^3 + 5\theta^2 + 8\theta + 4}{(\theta^2 + 2\theta + 2)^2} d\theta$$

28.
$$\int \frac{\theta^4 - 4\theta^3 + 2\theta^2 - 3\theta + 1}{(\theta^2 + 1)^3} d\theta$$

Improper Fractions

In Exercises 29–34, perform long division on the integrand, write the proper fraction as a sum of partial fractions, and then evaluate the integral.

29.
$$\int \frac{2x^3 - 2x^2 + 1}{x^2 - x} dx$$

30.
$$\int \frac{x^4}{x^2 - 1} dx$$

31.
$$\int \frac{9x^3 - 3x + 1}{x^3 - x^2} dx$$

32.
$$\int \frac{16x^3}{4x^2 - 4x + 1} dx$$

33.
$$\int \frac{y^4 + y^2 - 1}{y^3 + y} dy$$

34.
$$\int \frac{2y^4}{y^3 - y^2 + y - 1} dy$$

Evaluating Integrals

Evaluate the integrals in Exercises 35–40.

35. $\int \frac{e^t dt}{e^{2t} + 3e^t + 2}$ 36. $\int \frac{e^{4t} + 2e^{2t} - e^t}{e^{2t} + 1} dt$
37. $\int \frac{\cos y dy}{\sin^2 y + \sin y - 6}$ 38. $\int \frac{\sin \theta d\theta}{\cos^2 \theta + \cos \theta - 2}$
39. $\int \frac{(x-2)^2 \tan^{-1}(2x) - 12x^3 - 3x}{(4x^2 + 1)(x-2)^2} dx$
40. $\int \frac{(x+1)^2 \tan^{-1}(3x) + 9x^3 + x}{(9x^2 + 1)(x+1)^2} dx$



Initial Value Problems

Solve the initial value problems in Exercises 41–44 for x as a function of t .

41. $(t^2 - 3t + 2) \frac{dx}{dt} = 1 \quad (t > 2), \quad x(3) = 0$
42. $(3t^4 + 4t^2 + 1) \frac{dx}{dt} = 2\sqrt{3}, \quad x(1) = -\pi\sqrt{3}/4$
43. $(t^2 + 2t) \frac{dx}{dt} = 2x + 2 \quad (t, x > 0), \quad x(1) = 1$
44. $(t + 1) \frac{dx}{dt} = x^2 + 1 \quad (t > -1), \quad x(0) = \pi/4$


Applications and Examples

In Exercises 45 and 46, find the volume of the solid generated by revolving the shaded region about the indicated axis.

45. The x -axis46. The y -axis

- T** 47. Find, to two decimal places, the x -coordinate of the centroid of the region in the first quadrant bounded by the x -axis, the curve $y = \tan^{-1} x$, and the line $x = \sqrt{3}$.

- T** 48. Find the x -coordinate of the centroid of this region to two decimal places.



- T** 49. **Social diffusion** Sociologists sometimes use the phrase “social diffusion” to describe the way information spreads through a population. The information might be a rumor, a cultural fad, or news about a technical innovation. In a sufficiently large population, the number of people x who have the information is treated as a differentiable function of time t , and the rate of diffusion, dx/dt , is assumed to be proportional to the number of people who have the information times the number of people who do not. This leads to the equation

$$\frac{dx}{dt} = kx(N - x),$$

where N is the number of people in the population.Suppose t is in days, $k = 1/250$, and two people start a rumor at time $t = 0$ in a population of $N = 1000$ people.

- Find x as a function of t .
 - When will half the population have heard the rumor? (This is when the rumor will be spreading the fastest.)
- T** 50. **Second-order chemical reactions** Many chemical reactions are the result of the interaction of two molecules that undergo a change to produce a new product. The rate of the reaction typically depends on the concentrations of the two kinds of molecules. If a is the amount of substance A and b is the amount of substance B at time $t = 0$, and if x is the amount of product at time t , then the rate of formation of x may be given by the differential equation

$$\frac{dx}{dt} = k(a - x)(b - x),$$

or

$$\frac{1}{(a - x)(b - x)} \frac{dx}{dt} = k,$$

where k is a constant for the reaction. Integrate both sides of this equation to obtain a relation between x and t (a) if $a = b$, and (b) if $a \neq b$. Assume in each case that $x = 0$ when $t = 0$.

51. **An integral connecting π to the approximation $22/7$**

- Evaluate $\int_0^1 \frac{x^4(x-1)^4}{x^2+1} dx$.
- How good is the approximation $\pi \approx 22/7$? Find out by expressing $\left(\frac{22}{7} - \pi\right)$ as a percentage of π .

- c. Graph the function $y = \frac{x^4(x-1)^4}{x^2+1}$ for $0 \leq x \leq 1$. Experiment with the range on the y -axis set between 0 and 1, then between 0 and 0.5, and then decreasing the range until the graph can be seen. What do you conclude about the area under the curve?

52. Find the second-degree polynomial $P(x)$ such that $P(0) = 1$, $P'(0) = 0$, and

$$\int \frac{P(x)}{x^3(x-1)^2} dx$$

is a rational function.