



INDIAN INSTITUTE OF SCIENCE

STOCHASTIC HYDROLOGY

Lecture -20

Course Instructor : Prof. P. P. MUJUMDAR

Department of Civil Engg., IISc.

Summary of the previous lecture

- Case study -3: Monthly streamflows ate KRS reservoir
 - Plots of Time series, Correlogram, Partial Autocorrelation function and Power spectrum
 - Standardization to remove periodicities
 - Candidate ARMA models : (contiguous and non-contiguous)
 - Log Likelihood
 - Mean square error

CASE STUDIES - ARMA MODELS

Case study – 3

Monthly Stream flow (cum/sec) Statistics(1934 -1974) for Cauvery River at Krishna Raja Sagar Reservoir is considered in the case study.

- Time series of the data, auto correlation function, partial auto correlation function and the power spectrum are plotted.
- The series indicates presence of periodicities.
- The series is standardized to remove the periodicities.

Case study – 3 (Contd.)

- Standardized series is considered for fitting the ARMA models
- Total length of the data set $N = 480$
- Half the data set (240 values) is used to construct the model and other half is used for validation.
- Both contiguous and non-contiguous models are studied
- Non-contiguous models consider the most significant AR and MA terms leaving out the intermediate terms

Case study – 3 (Contd.)

Contiguous models:

$$L_i = -\frac{N}{2} \ln(\sigma_i) - n_i$$

Sl. No	Model	Likelihood values
1		29.33
2	ARMA(2,0)	28.91
3	ARMA(3,0)	28.96
4	ARMA(4,0)	31.63
5	ARMA(5,0)	30.71
6	ARMA(6,0)	29.90
	ARMA(1,1)	30.58
8	ARMA(1,2)	29.83
9	ARMA(2,1)	29.83
10	ARMA(2,2)	28.80
11	ARMA(3,1)	29.45

Case study – 3 (Contd.)

Non-contiguous models*:

Sl. No	Model	Likelihood values
1	ARMA(2,0)	28.52
2	ARMA(3,0)	28.12
3	ARMA(4,0)	28.21
4	ARMA(5,0)	30.85
5	ARMA(6,0)	29.84
6	ARMA(7,0)	29.12
7	ARMA(2,2)	29.81
8	ARMA(2,3)	28.82
9	ARMA(3,2)	28.48
10	ARMA(3,3)	28.06
11	ARMA(4,2)	28.65

*: The last AR and MA terms correspond to the 12th lag

Case study – 3 (Contd.)

- For this time series, the likelihood values for
 - contiguous model = 31.63
 - non-contiguous model = 30.85
- Hence contiguous ARMA(4,0) can be used.
- The parameters for the selected model are as follows

$$\phi_1 = 0.2137$$

$$\phi_2 = 0.0398$$

$$\phi_3 = 0.054$$

$$\phi_4 = 0.1762$$

$$\text{Constant} = -0.0157$$

Case study – 3 (Contd.)

Forecasting Models

Contiguous models:

Sl. No	Model	Mean square error values
1	ARMA(1,0)	0.97
2	ARMA(2,0)	1.92
3	ARMA(3,0)	2.87
4	ARMA(4,0)	3.82
5	ARMA(5,0)	4.78
6	ARMA(6,0)	5.74
7	ARMA(1,1)	2.49
8	ARMA(1,2)	2.17
9	ARMA(2,1)	3.44
10	ARMA(2,2)	4.29
11	ARMA(3,1)	1.89

Case study – 3 (Contd.)

Non-contiguous models:

Sl. No	Model	Mean square error values
1	ARMA(2,0)	0.96
2	ARMA(3,0)	1.89
3	ARMA(4,0)	2.84
4	ARMA(5,0)	3.79
5	ARMA(6,0)	4.74
6	ARMA(7,0)	5.7
7	ARMA(2,2)	2.42
8	ARMA(2,3)	1.99
9	ARMA(3,2)	2.52
10	ARMA(3,3)	1.15
11	ARMA(4,2)	1.71

Case study – 3 (Contd.)

- The simplest model AR(1) results in the least value of the MSE
- For one step forecasting, quite often the simplest model is appropriate
- Also as the number of parameters increases, the MSE increases which is contrary to the common belief that models with large number of parameters give better forecasts.
- AR(1) model is recommended for forecasting the series and the parameters are as follows
 $\phi_1 = 0.2557$ and $C = -0.009$

Case study – 3 (Contd.)

- Validation tests on the residual series
 - Significance of residual mean
 - Significance of periodicities
 - Cumulative periodogram test or Bartlett's test
 - White noise test

- Residuals,
$$e_t = X_t - \left(\underbrace{\sum_{j=1}^{m_1} \phi_j X_{t-j} + \sum_{j=1}^{m_2} \theta_j e_{t-j}}_{\text{Simulated from the model}} + C \right)$$

Residual

Data

Simulated from the model

Case study – 3 (Contd.)

Significance of residual mean:

Sl. No	Model	$\eta(e)$	$t_{0.95}(239)$
1	ARMA(1,0)	0.002	1.645
2	ARMA(2,0)	0.006	1.645
3	ARMA(3,0)	0.008	1.645
4	ARMA(4,0)	0.025	1.645
5	ARMA(5,0)	0.023	1.645
6	ARMA(6,0)	0.018	1.645
7	ARMA(1,1)	0.033	1.645
8	ARMA(1,2)	0.104	1.645
9	ARMA(2,1)	0.106	1.645
10	ARMA(2,2)	0.028	1.645


$$\eta(e) = \frac{N^{1/2}e}{\hat{\rho}^{1/2}}$$

$\eta(e) \leq t(0.95, 240-1)$;
All models pass the test

Case study – 3 (Contd.)

Significance of periodicities:

$$\eta(e) = \frac{\gamma_k^2 (N - 2)}{4\hat{\rho}_1}$$



$$\gamma_k^2 = \alpha_k^2 + \beta_k^2$$

$$\hat{\rho}_1 = \frac{1}{N} \left[\sum_{t=1}^N \left\{ e_t - \hat{\alpha} \cos(\omega_k t) - \hat{\beta} \sin(\omega_k t) \right\}^2 \right]$$

$$\alpha_k = \frac{2}{N} \sum_{t=1}^n e_t \cos(\omega_k t)$$

Case study – 3 (Contd.)

$$\beta_k = \frac{2}{N} \sum_{t=1}^n e_t \sin(\omega_k t)$$

$2\pi/\omega_k$ is the periodicity for which test is being carried out.

$\eta(e) \leq F_\alpha(2, N-2)$ – Model passes the test

Case study – 3 (Contd.)

Significance of periodicities:

Sl. No	Model	η Value for the periodicity				$F_{0.95}(2,238)$
		1 st	2 nd	3 rd	4 th	
1	ARMA(1,0)	0.527	1.092	0.364	0.065	3.00
2	ARMA(2,0)	1.027	2.458	0.813	0.129	3.00
3	ARMA(3,0)	1.705	4.319	1.096	0.160	3.00
4	ARMA(4,0)	3.228	6.078	0.948	0.277	3.00
5	ARMA(5,0)	3.769	7.805	1.149	0.345	3.00
6	ARMA(6,0)	4.19	10.13	1.262	0.441	3.00
7	ARMA(1,1)	4.737	10.09	2.668	0.392	3.00
8	ARMA(1,2)	6.786	10.67	2.621	0.372	
9	ARMA(2,1)	7.704	12.12	2.976	0.422	3.00
10	ARMA(2,2)	6.857	13.22	3.718	0.597	3.00


Case study – 3 (Contd.)

Significance of periodicities by Bartlett's test :
(Cumulative periodogram test)

$$\gamma_k^2 = \left\{ \frac{2}{N} \sum_{t=1}^N e_t \cos(\omega_k t) \right\}^2 + \left\{ \frac{2}{N} \sum_{t=1}^N e_t \sin(\omega_k t) \right\}^2$$
$$g_k = \frac{\sum_{j=1}^k \gamma_j^2}{\sum_{k=1}^{N/2} \gamma_k^2} \quad k = 1, 2, \dots, N/2$$

Case study – 3 (Contd.)

Cumulative periodogram for the original series without standardizing




Case study – 3 (Contd.)

- The confidence limits ($1.35/(N/2)^{1/2} = 0.087$) are plotted for 95% confidence. ($N = 480$ in this case)
- Most part of the cumulative periodogram lies outside the significance bands confirming the presence of periodicity in the data.
- for $k=40$, a spike in the graph is seen indicating the significant periodicity
- This ‘ k ’ corresponds to a periodicity of 12 months ($480/40$)
- $k=80$, corresponds to a periodicity of 6 months

Case study – 3 (Contd.)

Cumulative periodogram for the residual series of ARMA(4, 0) model



Case study – 3 (Contd.)

- The confidence limits ($\pm 1.35/(N/2)^{1/2} = \pm 0.123$) are plotted for 95% confidence. ($N = 240$, in this case)
- Cumulative periodogram lies within the significance bands confirming that no significant periodicity present in the residual series.
- The model passes the test.

Whittle's test for white noise

White noise test (Whittle's test):

- This test is carried out to test the absence of correlation in the series.
- The covariance r_k at lag k of the error series $\{e_t\}$

$$r_k = \frac{1}{N-k} \sum_{j=k+1}^N e_j e_{j-k} \quad k = 0, 1, 2, \dots, k_{\max}$$

- The value of k_{\max} is normally chosen as $0.15N$

Ref: Kashyap R.L. and Ramachandra Rao.A, "Dynamic stochastic models from empirical data", Academic press, New York, 1976

Whittle's test for white noise

- The covariance matrix is

$$\Gamma_{n1} = \begin{bmatrix} r_0 & r_1 & r_2 & \cdot & \cdot & r_{k_{\max}} \\ r_1 & r_0 & r_1 & \cdot & \cdot & r_{k_{\max}-1} \\ r_2 & & & & & \\ \cdot & & & & & \\ \cdot & & & & & \\ r_{k_{\max}} & r_{k_{\max}-1} & & & r_0 & \end{bmatrix} k_{\max} \times k_{\max}$$

Whittle's test for white noise

- A statistic $\eta(e)$ is defined as

$$\eta(e) = \frac{N}{n_1 - 1} \left(\frac{\hat{\rho}_0}{\hat{\rho}_1} - 1 \right) \quad n_1 = k_{\max}$$

Where $\hat{\rho}_0$ is the lag zero correlation =1, and

$$\hat{\rho}_1 = \frac{\det \Gamma_{n_1}}{\det \Gamma_{n_1-1}}$$

The matrix Γ_{n_1-1} is constructed by eliminating the last row and the last column from the Γ_{n_1} matrix.

Whittle's test for white noise

- The statistic $\eta(e)$ is approximately distributed as $F_\alpha(n_1, N-n_1)$, where α is the significance level at which the test is being carried out.
- If the value of $\eta(e) \leq F_\alpha(n_1, N-n_1)$, then the residual series is uncorrelated.

$F_{\alpha}(n, N-n)$

Case study – 3 (Contd.)

Whittle's test - white noise

$$\eta(e) = \frac{N}{n_1-1} \left(\frac{\hat{\rho}_0}{\hat{\rho}_1} - 1 \right)$$

Model \ n1	n1 = 73	n1 = 49	n1 = 25
Model	η	η	η
ARMA(1,0)	0.642	0.917	0.891
ARMA(2,0)	0.628	0.898	0.861
ARMA(3,0)	0.606	0.868	0.791
ARMA(4,0)	0.528	0.743	0.516
ARMA(5,0)	0.526	0.739	0.516
ARMA(6,0)	0.522	0.728	0.493
ARMA(1,1)	0.595	0.854	0.755
ARMA(1,2)	0.851	1.256	1.581
ARMA(2,1)	0.851	1.256	1.581
ARMA(2,2)	0.589	0.845	0.737

 model fails

Case study – 3 (Contd.)

$$\eta(e) = (N - n_1) \sum_{k=1}^{n_1} \left(\frac{r_k}{r_0} \right)^2$$


Portmanteau test for white noise:

$\chi^2_{0.95}(k_{\max})$	kmax = 48	kmax = 36	kmax = 24	kmax = 12
Model	η	η	η	η
ARMA(1,0)	31.44	33.41	23.02	14.8
ARMA(2,0)	32.03	34.03	24.47	15.17
ARMA(3,0)	30.17	32.05	21.61	13.12
ARMA(4,0)	20.22	21.49	11.85	4.31
ARMA(5,0)	19.84	21.08	11.75	4.14
ARMA(6,0)	19.64	20.87	11.48	3.79
ARMA(1,1)	29.89	31.76	22.24	12.76
ARMA(1,2)	55.88	59.38	48.37	39.85
ARMA(2,1)	55.88	59.38	48.37	38.85
ARMA(2,2)	28.62	30.41	20.39	11.25

 model fails


Case study – 4

Monthly Stream flow (1928-1964) of a river (37 years of monthly data);




Case study – 4 (Contd.)

- Correlogram




Case study – 4 (Contd.)

- Partial Auto Correlation function



Case study – 4 (Contd.)

- Power Spectrum



Case study – 4 (Contd.)

Model name	constant	ϕ_1	ϕ_2	ϕ_3	ϕ_4	ϕ_5	ϕ_6	ϕ_7	ϕ_8	θ_1	θ_2
ARMA(1,0)	-0.097	0.667									
ARMA(2,0)	0.049	0.042	0.044								
ARMA(3,0)	-0.111	0.767	-0.148	-0.003							
ARMA(4,0)	0.052	0.042	0.058	0.063	0.044						
ARMA(5,0)	0.055	0.042	0.058	0.063	0.048	0.038					
ARMA(6,0)	-0.124	0.764	-0.155	0.034	-0.029	-0.023	-0.021				
ARMA(7,0)	0.056	0.043	0.062	0.065	0.048	0.058	0.075	0.065			
ARMA(8,0)	0.054	0.042	0.060	0.063	0.048	0.058	0.078	0.088	0.061		
ARMA(1,1)	-0.131	0.551								0.216	
ARMA(2,1)	-0.104	0.848	-0.204							-0.083	
ARMA(3,1)	-0.155	0.351	0.165	-0.055						0.418	
ARMA(4,1)	-0.083	1.083	-0.400	0.091	-0.060					-0.318	
ARMA(1,2)	-0.139	0.526								0.241	0.025
ARMA(2,2)	378	1980	1160							1960	461
ARMA(0,1)	-0.298									0.594	
ARMA(0,2)	-0.297									0.736	0.281

Case study – 4 (Contd.)

Sl. No	Model	Mean Square Error	Likelihood value
1	ARMA(1,0)	0.65	93.33
2	ARMA(2,0)	0.63	97.24
3	ARMA(3,0)	0.63	96.44
4	ARMA(4,0)	0.63	96.14
5	ARMA(5,0)	0.63	95.50
6	ARMA(6,0)	0.63	94.66
7	ARMA(7,0)	0.63	93.80
8	ARMA(8,0)	0.60	101.47
9	ARMA(1,1)	0.63	97.11
10	ARMA(2,1)	0.63	96.25
11	ARMA(3,1)	0.63	95.39
12	ARMA(4,1)	0.63	95.39
13	ARMA(1,2)	0.63	96.16
14	ARMA(2,2)	0.63	95.13
15	ARMA(0,1)	0.73	67.74
16	ARMA(0,2)	0.66	89.28

Case study – 4 (Contd.)

Significance of residual mean:

$$\eta(e) = \frac{N^{1/2}\bar{e}}{\hat{\rho}^{1/2}}$$

\bar{e} is the estimate of the residual mean


$\hat{\rho}$ is the estimate of the residual variance

- All models pass the test

Model	Test t	
	$\eta(e)$	$t(\alpha, N-1)$
ARMA(1,0)	1.78672E-05	1.645
ARMA(2,0)	6.0233E-06	1.645
ARMA(3,0)	4.82085E-05	1.645
ARMA(4,0)	-3.01791E-05	1.645
ARMA(5,0)	6.84076E-16	1.645
ARMA(6,0)	3.0215E-05	1.645
ARMA(7,0)	-6.04496E-06	1.645
ARMA(8,0)	5.54991E-05	1.645
ARMA(1,1)	-0.001132046	1.645
ARMA(2,1)	-0.002650292	1.645
ARMA(3,1)	-0.022776166	1.645
ARMA(4,1)	0.000410668	1.645
ARMA(1,2)	-0.000837092	1.645
ARMA(2,2)	0.002631505	1.645
ARMA(0,1)	0.022950466	1.645
ARMA(0,2)	0.019847826	1.645


Case study – 5

Sakleshpur Annual Rainfall Data (1901-2002)




Case study – 5 (Contd.)

Correlogram



Case study – 5 (Contd.)

PAC function



Case study – 5 (Contd.)

Model	Likelihood
AR(1)	9.078037
AR(2)	8.562427
AR(3)	8.646781
AR(4)	9.691461
AR(5)	9.821681
AR(6)	9.436822
ARMA(1,1)	8.341717
ARMA(1,2)	8.217627
ARMA(2,1)	7.715415
ARMA(2,2)	5.278434
ARMA(3,1)	6.316174
ARMA(3,2)	6.390390

Case study – 5 (Contd.)

- ARMA(5,0) is selected with highest likelihood value
- The parameters for the selected model are as follows

$$\phi_1 = 0.40499$$

$$\phi_2 = 0.15223$$

$$\phi_3 = -0.02427$$

$$\phi_4 = -0.2222$$

$$\phi_5 = 0.083435$$

$$\text{Constant} = -0.000664$$

Case study – 5 (Contd.)

- Significance of residual mean

Model	$\eta(e)$	$t_{0.95}(N)$
	0.000005	1.6601

Case study – 5 (Contd.)

Significance of periodicities:

Periodicity	η	$F_{0.95}(2,239)$
1 st	0.000	3.085
2 nd	0.00432	3.085
3 rd	0.0168	3.085
4 th	0.0698	3.085
5 th	0.000006	3.085
6 th	0.117	3.085

Case study – 5 (Contd.)

- Whittle's white noise test:

Model	η	$F_{0.95}(2, N-2)$
ARMA(5,0)	0.163	1.783

Case study – 5 (Contd.)

Model	MSE
AR(1)	1.180837
AR(2)	1.169667
AR(3)	1.182210
AR(4)	1.168724
AR(5)	1.254929
AR(6)	1.289385
ARMA(1,1)	1.171668
ARMA(1,2)	1.156298
ARMA(2,1)	1.183397
ARMA(2,2)	1.256068
ARMA(3,1)	1.195626
ARMA(3,2)	27.466087

Case study – 5 (Contd.)

- ARMA(1, 2) is selected with least MSE value for one step forecasting
- The parameters for the selected model are as follows

$$\phi_1 = 0.35271$$

$$\theta_1 = 0.017124$$

$$\theta_2 = -0.216745$$

$$\text{Constant} = -0.009267$$

Case study – 5 (Contd.)

- Significance of residual mean

Model	$\eta(e)$	$t_{0.95}(N)$
ARMA(1, 2)	-0.0026	1.6601

Case study – 5 (Contd.)

Significance of periodicities:

Periodicity	η	$F_{0.95}(2,239)$
1 st	0.000	3.085
2 nd	0.0006	3.085
3 rd	0.0493	3.085
4 th	0.0687	3.085
5 th	0.0003	3.085
6 th	0.0719	3.085

Case study – 5 (Contd.)

- Whittle's white noise test:

Model	η	$F_{0.95}(2, N-2)$
ARMA(1, 2)	0.3605	1.783

Markov Chains

Markov Chains

- A Markov chain is a stochastic process with the property that value of process X_t at time t depends on its value at time $t-1$ and not on the sequence of other values ($X_{t-2}, X_{t-3}, \dots, X_0$) that the process passed through in arriving at X_{t-1} .

$$P[X_t/X_{t-1}, X_{t-2}, \dots, X_0] = P[X_t/X_{t-1}]$$

↗ Single step Markov chain

Markov Chains

$$P[X_t = a_j / X_{t-1} = a_i]$$

- This conditional probability gives the probability at time t will be in state ‘j’ , given that the process was in state ‘i’ at time t-1.
- The conditional probability is independent of the states occupied prior to t-1.
- For example, if X_{t-1} is a dry day, we would be interested in the probability that X_t is a dry day or a wet day.
- This probability is commonly called as transition probability

Markov Chains

$$P[X_t = a_j | X_{t-1} = a_i] = P_{ij}^t$$

- Usually written as P_{ij}^t indicating the probability of a step from a_i to a_j at time 't'.
- If P_{ij} is independent of time, then the Markov chain is said to be homogeneous.

$$\text{i.e., } P_{ij}^t = P_{ij}^{t+\tau} \quad \forall \quad t \text{ and } \tau$$

the transition probabilities remain same across time

Markov Chains

Transition Probability Matrix(TPM):

$$P = \begin{bmatrix} & \begin{matrix} t+1 \rightarrow & 1 & 2 & 3 & \cdot & \cdot & m \end{matrix} \\ \begin{matrix} t \\ \downarrow \\ 1 \\ 2 \\ 3 \\ \cdot \\ \cdot \\ m \end{matrix} & \begin{bmatrix} P_{11} & P_{12} & P_{13} & \cdot & \cdot & P_{1m} \\ P_{21} & P_{22} & P_{23} & \cdot & \cdot & P_{2m} \\ P_{31} \\ \cdot \\ \cdot \\ P_{m1} & P_{m2} & & & & P_{mm} \end{bmatrix} \end{bmatrix}_{m \times m}$$

Markov Chains

$$\sum_{j=1}^m P_{ij} = 1 \quad \forall i$$

- Elements in any row of TPM sum to unity
- TPM can be estimated from observed data by enumerating the number of times the observed data went from state ‘i’ to ‘j’
- $P_j^{(n)}$ is the probability of being in state ‘j’ in time step ‘n’.

Markov Chains

- $p_j^{(0)}$ is the probability of being in state ‘j’ in period $t = 0$.

$$p^{(0)} = \begin{bmatrix} p_1^{(0)} & p_2^{(0)} & \dots & p_m^{(0)} \end{bmatrix}_{1 \times m} \quad \dots \text{Probability vector at time 0}$$

$$p^{(n)} = \begin{bmatrix} p_1^{(n)} & p_2^{(n)} & \dots & p_m^{(n)} \end{bmatrix}_{1 \times m} \quad \dots \text{Probability vector at time 'n'}$$

- If $p^{(0)}$ is given and TPM is given

$$p^{(1)} = p^{(0)} \times P$$

Markov Chains

$$p^{(1)} = \begin{bmatrix} p_1^{(0)} & p_2^{(0)} & \dots & p_m^{(0)} \end{bmatrix} \begin{bmatrix} P_{11} & P_{12} & P_{13} & \dots & P_{1m} \\ P_{21} & P_{22} & P_{23} & \dots & P_{2m} \\ P_{31} \\ \vdots \\ P_{m1} & P_{m2} & & & P_{mm} \end{bmatrix}$$

$$= p_1^{(0)}P_{11} + p_2^{(0)}P_{21} + \dots + p_m^{(0)}P_{m1} \quad \dots \text{Probability of going to state 1}$$

$$= p_1^{(0)}P_{12} + p_2^{(0)}P_{21} + \dots + p_m^{(0)}P_{m2} \quad \dots \text{Probability of going to state 2}$$

And so on...

Markov Chains

Therefore

$$p^{(1)} = \begin{bmatrix} p_1^{(1)} & p_2^{(1)} & \dots & p_m^{(1)} \end{bmatrix}_{1 \times m}$$

$$\begin{aligned} p^{(2)} &= p^{(1)} \times P \\ &= p^{(0)} \times P \times P \\ &= p^{(0)} \times P^2 \end{aligned}$$

In general,

$$p^{(n)} = p^{(0)} \times P^n$$

Markov Chains

- As the process advances in time, $p_j^{(n)}$ becomes less dependent on $p^{(0)}$
- The probability of being in state ‘j’ after a large number of time steps becomes independent of the initial state of the process.
- The process reaches a steady state at large n

$$p = p \times P^n$$

- As the process reaches steady state, the probability vector remains constant

Example – 1

Consider the TPM for a 2-state first order homogeneous Markov chain as

$$TPM = \begin{bmatrix} 0.7 & 0.3 \\ 0.4 & 0.6 \end{bmatrix}$$

State 1 is a non-rainy day and state 2 is a rainy day

Obtain the

1. probability of day 1 is non-rainfall day / day 0 is rainfall day
2. probability of day 2 is rainfall day / day 0 is non-rainfall day
3. probability of day 100 is rainfall day / day 0 is non-rainfall day

Example – 1 (contd.)

1. probability of day 1 is non-rainfall day / day 0 is rainfall day

$$TPM = \begin{matrix} & \text{No rain} & \text{rain} \\ \text{No rain} & [0.7 & 0.3] \\ \text{rain} & [0.4 & 0.6] \end{matrix}$$

The probability is 0.4

2. probability of day 2 is rainfall day / day 0 is non-rainfall day

$$p^{(2)} = p^{(0)} \times P^2$$

Example – 1 (contd.)

$$\begin{aligned} p^{(2)} &= [0.7 \quad 0.3] \begin{bmatrix} 0.7 & 0.3 \\ 0.4 & 0.6 \end{bmatrix} \\ &= [0.61 \quad 0.39] \end{aligned}$$

The probability is 0.39

3. probability of day 100 is rainfall day / day 0 is non-rainfall day

$$p^{(n)} = p^{(0)} \times P^n$$

Example – 1 (contd.)

$$P^2 = P \times P$$

$$= \begin{bmatrix} 0.7 & 0.3 \\ 0.4 & 0.6 \end{bmatrix} \begin{bmatrix} 0.7 & 0.3 \\ 0.4 & 0.6 \end{bmatrix} = \begin{bmatrix} 0.61 & 0.39 \\ 0.52 & 0.48 \end{bmatrix}$$

$$P^4 = P^2 \times P^2 = \begin{bmatrix} 0.5749 & 0.4251 \\ 0.5668 & 0.4332 \end{bmatrix}$$

$$P^8 = P^4 \times P^4 = \begin{bmatrix} 0.5715 & 0.4285 \\ 0.5714 & 0.4286 \end{bmatrix}$$

$$P^{16} = P^8 \times P^8 = \begin{bmatrix} 0.5714 & 0.4286 \\ 0.5714 & 0.4286 \end{bmatrix}$$

Example – 1 (contd.)

Steady state probability

$$p = [0.5714 \quad 0.4286]$$

For steady state,

$$p = p \times P^n$$

$$\begin{aligned} &= [0.5714 \quad 0.4286] \begin{bmatrix} 0.5714 & 0.4286 \\ 0.5714 & 0.4286 \end{bmatrix} \\ &= [0.5714 \quad 0.4286] \end{aligned}$$